

Deletion in the FMR1 Gene in a Fragile-X Male

Arto Mannermaa, Leena Pulkkinen, Eliisa Kajanoja, Markku Ryyänen, and Seppo Saarikoski

Department of Obstetrics and Gynecology, Division of Clinical Genetics, Kuopio University Hospital, Finland

The pathogenesis of Fragile-X syndrome is a consequence of absence of the FMR1 gene product associated with expansion of the CGG repeat and abnormal methylation of this and a CpG island 250 bp proximal to the CGG repeat located at exon 1 in the FMR1 gene. While this is usually the case, some suspected Fragile-X syndrome patients have been described with a mutation other than CGG expansion. We describe here an affected Fragile-X male, who was found to be mosaic of a full mutation of the CGG expansion and a deletion in the FMR1 gene. The patient's phenotype is probably mainly due to the effect of the full mutation of the repeat sequence. Thus, the influence of the deletion is difficult to evaluate.

© 1996 Wiley-Liss, Inc.

KEY WORDS: mental retardation, deletion, mosaicism

INTRODUCTION

Fragile-X syndrome is one of the most common forms of mental retardation, with a frequency of 1 to 1,500 males and 1 to 2,500 females [Webb et al., 1986], and manifestations of severe mental retardation, developmental delay, and such physical characteristics as large ears, long face with a prominent jaw, and macroorchidism. This syndrome is inherited as an X-linked dominant disorder with reduced penetrance. Fragile-X syndrome is associated with a folate-sensitive fragile site, FRAXA, at Xq27.3 [Lubs, 1969].

The pathogenesis of Fragile-X syndrome is a consequence of absence of the FMR1 gene product associated with expansion and abnormal methylation of the CGG repeat located at exon 1 in the FMR1 gene [Kremer et al., 1991; Oberlé et al., 1991; Verkerk et al., 1991; Yu et al., 1991]. While this is usually the case, some

suspected Fragile-X syndrome patients have been described with a mutation other than CGG expansion. Large deletions involving removal of the whole FMR1 gene have been reported [Gedeon et al., 1992; Tarleton et al., 1993; Gu et al., 1994; Quan et al., 1995] but shorter deletions have also been found, where absence of FMR1-expression can be explained by loss of sequences proximal to the FMR1 gene and/or sequences flanking and including the CGG repeat region [Wohrle et al., 1992; Meijer et al., 1994; Trottier et al., 1994; Hirst et al., 1995]. In addition, de Boule et al. [1993] have described a point mutation resulting in a 367^{Le}→367^{Asn} substitution in the FMR1 gene and leading to the Fragile-X phenotype.

We describe here a detailed sequence analysis of the FMR1 gene of an affected Fragile-X male, who was found to be mosaic for a full mutation of the CGG expansion and a deletion in the 2.8 kb *EcoRI-EagI* fragment recognized by the StB12.3 probe.

MATERIALS AND METHODS

Case Presentation

The index case is a 52-year-old Finnish man with moderate mental retardation, with no ability to read or write, now attending a sheltered workshop. He has a classic Fragile-X phenotype with obesity. Of the family, only a phenotypically healthy sister is alive.

DNA Analysis

The probe StB12.3 [Oberlé et al., 1991] was used for Southern blot analysis of the FMR1 gene. PCR analysis of the deleted area was carried out as described by Chong et al. [1994] with the primers 5'-GGAACAGCG-TTGATCACGTGACGTGGTTTC-3' and 5'-GGGGCCT-GCCCTAGSGCCAAGTACCTTGT-3'. The PCR product was subjected to direct sequencing utilizing a USB PCR product sequencing kit, with ³⁵S-dATP as an isotope, according to the manufacturer's instructions.

RESULTS AND DISCUSSION

Although most Fragile-X patients have an expanded CGG repeat sequence in the FMR1 gene, it is now becoming evident that other molecular mechanisms also exist leading to an affected phenotype. A full mutation with over 200 CGG repeats represses FMR1 transcription [Pieretti et al., 1991; Verheij et al., 1993] and results in reduction of the amount of translated FMR1 protein. In this study, analysis of the index case demon-

Received for publication September 13, 1995; revision received December 21, 1995.

Address reprint requests to Arto Mannermaa, Department of Obstetrics and Gynecology, Division of Clinical Genetics, Kuopio University Hospital, Building 2 D, FIN-70210, Finland.




Fig. 1. Southern blot showing the CGG repeat of the FMR1 of a fragile X male (II-2) and his relatives. DNA obtained from peripheral blood was digested with *EcoRI* and *EagI* and hybridized with StB12.3. The specimen of the proband shows a mosaic situation with a smear of full mutation size and an abnormally small fragment.

strated a mosaic banding pattern with a full methylated mutation in the CGG region, and a shortened *EcoRI-EagI* fragment compared with the 2.8 kb normal sized fragment (Fig. 1).

To further assess the size and location of the deletion in the FMR1 gene, PCR-based analysis (Fig. 2a) and direct sequencing was utilized. A deletion spanning the entire CGG-repeat and flanking sequences of 31 bp at the 5'-side and 61 bp at the 3'-side of the CGG-repeat within the untranslating region was found (Fig. 2b).

The patient described in this paper presents a rare type of mosaicism in Fragile-X males. Similar cases with both a full methylated mutation and a deletion in and adjacent to the CGG repeat sequence have previously been reported by de Graaf et al. [1995] and De Vries et al. [1993]. The deletion was found not to disrupt the coding sequence of the FMR1 gene. Although this alteration is located at the hotspot deletion region described by de Graaff et al. [1995], the influ-




Fig. 2. PCR amplification as described by Chong et al. [1994] and sequencing of the deleted area. **a:** Amplified PCR-products were sized in an agarose gel electrophoresis and stained with ethidium bromide. P, patient; C, control sample; MW *spI* digested pBR322 as a molecular weight marker. **b:** The sequenced PCR product from the untranslating region of the FMR1 gene of the proband demonstrates a deletion spanning the entire CGG repeat and flanking sequences of 31 bp and 61 bp.

ence of this deletion on FMR1 gene expression and protein synthesis is difficult to interpret, since the promoter region of the FMR1 gene is only partially described. Furthermore, the phenotypic effect of this deletion is probably obscured by the effect of the full mutation of the CGG repeat. It is also possible that the deletion is a consequence of the instability of the full mutation sized CGG repeat [de Graaff et al., 1995]. Additional information concerning the impact of the deletion described here could be gained through in vitro gene expression studies.

ACKNOWLEDGMENTS

We thank Dr. Salme Majuri for referring the patient and Mrs. Pirkko Jokela and Mrs. Inkeri Happonen for carrying out the Southern hybridization. We also thank the proband and his relatives for their cooperation.

REFERENCES

- Chong SS, Eichler EE, Nelson DL, Hughes MR (1994): Robust amplification and ethidium-visible detection of the fragile X syndrome CGG repeat using Pfu polymerase. *Am J Med Genet* 51:522-526.
- De Boule K, Verkerk AJMH, Reyniers E, Vits L, Hendrickx J, Van Roy B, Van den Bos F, de Graaff E, Oostra BA, Willems PJ (1993): A Point Mutation in the FMR-1 Gene Associated with Fragile-X Mental Retardation. *Nat Genet* 3:31-35.
- de Graaff E, Rouillard P, Willems PJ, Smits APT, Rousseau F, Oostra BA (1995): Hotspot for deletions in the CGG repeat region of FMR1 in fragile X patients. *Hum Mol Genet* 4:45-49.
- De Vries BBA, Fryns JP, Butler MG, Canziani F, Wesby-van Swaay E, van Hemel JO, Oostra BA, Halley DJJ, Niermeijer MF (1993): Clinical and Molecular Studies in Fragile-X Patients with a Prader-Willi-Like Phenotype. *J Med Genet* 30:761-766.
- Gedeon AK, Baker E, Robinson H, Partington MW, Gross B, Manca A, Korn B, Poustka A, Yu S, Sutherland GR, Mulley JC (1992): Fragile-X Syndrome Without CCG Amplification Has an FMR1 Deletion. *Nat Genet* 1:341-344.
- Gu YH, Lugenbeel KA, Vockley JG, Grody WW, Nelson DL (1994): A de novo deletion in FMR1 in a patient with developmental delay. *Hum Mol Genet* 3:1705-1706.
- Hirst M, Grewal P, Flannery A, Slatter R, Maher E, Barton D, Fryns JP, Davies K (1995): Two new cases of FMR1 deletion associated with mental impairment. *Am J Hum Genet* 56:67-74.
- Kremer EJ, Pritchard M, Lynch M, Yu S, Holman K, Baker E, Warren ST, Schlessinger D, Sutherland GR, Richards RI (1991): Mapping of DNA instability at the fragile X to a trinucleotide repeat sequence p(CGG)n. *Science* 252:1711-1714.
- Lubs HA (1969): A marker X chromosome. *Am J Hum Genet* 21:231-244.
- Meijer H, de Graaff E, Merckx DML, Jongbloed RJE, de Die-Smulders CEM, Engelen JJM, Fryns JP, Curfs PMG, Oostra BA (1994): A Deletion of 1.6 kb Proximal to the Cgg Repeat of the FMR1 Gene Causes the Clinical Phenotype of the Fragile X Syndrome. *Hum Mol Genet* 3:615-620.
- Oberlé L, Rousseau F, Heitz D, Kretz C, Devys D, Hanauer A, Boué J, Bertheas MF, Mandel JL (1991): Instability of a 550-base pair DNA segment and abnormal methylation in fragile X syndrome. *Science* 252:1097-1102.
- Pieretti M, Zhang F, Fu YH, Warren ST, Oostra BA, Caskey CT, Nelson DL (1991): Absence of expression of the FMR-1 gene in fragile X syndrome. *Cell* 66:817-822.
- Quan F, Zonana J, Gunter K, Peterson KL, Magenis RE, Popovich BW (1995): An atypical case of fragile X syndrome caused by a deletion that includes the FMR1 gene. *Am J Hum Genet* 56:1042-1051.
- Tarleton J, Richie R, Schwartz C, Rao K, Aylsworth AS, Lachiewicz A (1993): An extensive de novo deletion removing FMR1 in a patient with mental retardation and the Fragile-X syndrome phenotype. *Hum Mol Genet* 2:1973-1974.
- Trottier Y, Imbert G, Poutska A, Fryns J-P, Mandel J-L (1994): Male with typical fragile X phenotype is deleted for part of the FMR1 gene and for about 10 kb of upstream region. *Am J Med Genet* 51:454-457.
- Verheij C, Bakker CE, de Graaff E, Keulemans J, Willemsen R, Verkerk AJMH, Galjaard H, Reuser AJJ, Hoogeveen AT, Oostra BA (1993): Characterization and localization of the FMR-1 gene product associated with Fragile-X syndrome. *Nature* 363:722-724.
- Verkerk AJMH, Pieretti M, Sutcliffe JS, Fu YH, Kuhl DPA, Pizzuti A, Reiner O, Richards S, Victoria MF, Zhang F, Eussen BE, van Ommen GJB, Blonden LAJ, Riggins GJ, Chastain JL, Kunst CB, Galjaard H, Caskey CT, Nelson DL, Oostra BA, Warren ST (1991): Identification of a gene (FMR-1) containing a CGG repeat coincident with a breakpoint cluster region exhibiting length variation in fragile X syndrome. *Cell* 65:905-914.
- Webb TP, Bunday SE, Thake AI, Todd J (1986): Population incidence and segregation ratios in the Martin-Bell syndrome. *Am J Med Genet* 23:573-580.
- Wohrle D, Kotzot D, Hirst MC, Manca A, Korn B, Schmidt A, Barbi G, Rott HD, Poutska A, Davies KE, Steinbach P (1992): A microdeletion of less than 250 kb, including the proximal part of the FMR1 gene and the fragile-X site, in a male with the clinical phenotype of fragile-X syndrome. *Am J Hum Genet* 51:299-306.
- Yu S, Pritchard M, Kremer E, Lynch M, Nancarrow J, Baker E, Holman K, Mulley JC, Warren ST, Schlessinger D, Sutherland GR, Richards RI (1991): Fragile X genotype characterized by an unstable region of DNA. *Science* 252:1179-1181.